

## ASK YOUR KIDS CARE FOR YOUR KIDS EMBRACE YOUR KIDS



See Reverse for Questions that Can Save a Life

Always ask questions 1 and 2.	Past	Month
1) Have you wished you were dead or wished you could go to sleep and not wake up?		
2) Have you actually had any thoughts about killing yourself?		
If <b>YES</b> to 2, ask questions 3, 4, 5 and 6. If <b>NO</b> to 2, skip to question 6.		
3) Have you been thinking about how you might do this?		
4) Have you had these thoughts and had some intention of acting on them?	High Risk	
5) Have you started to work out or worked out the details of how to kill yourself? Did you intend to carry out this plan?	High Risk	
Always Ask Question 6	Life- time	Past 3 Months
6) Have you done anything, started to do anything, or prepared to do anything to end your life?		High Risk
Examples: Collected pills, obtained a gun, gave away valuables, wrote a will or suicide note, took out pills but didn't swallow any, held a gun but changed your mind or it was grabbed from your hand, went to the roof but didn't jump; or actually took pills, tried to shoot yourself, cut yourself, tried to hang yourself, etc.		



Any YES indicates that someone should seek behavioral healthcare.

However, if the answer to 4, 5 or 6 is YES, seek immediate help: go to the ER, call 1-800-273-8255, text 741741 or call 911.

STAY WITH THEM until they can be evaluated.





Download Columbia Protocol app