CHICIDAL IDEATION								
SUICIDAL IDEATION								
Ask questions 1 and 2. If both are negative, proceed to "Suicidal Behavior" section. If the answer to question 2 is "yes", ask questions 3, 4 and 5. If the answer to question 1 and/or 2 is "yes", complete			Lifetime: Time He/She Felt		Past			
"Intensity of Ideation" section below.			uicidal	Mon	ıths			
1. Wish to be Dead								
Subject endorses thoughts about a wish to be dead or not alive anymore		Yes	No	Yes	No			
Have you wished you were dead or wished you could go to sleep and r	tot wake up?							
If yes, describe:								
2. Non-Specific Active Suicidal Thoughts			No	Yes	No			
General non-specific thoughts of wanting to end one's life/commit suicide (e.g., "I've thought about killing myself") without thoughts of ways to kill oneself/associated methods, intent, or plan during the assessment period.		Yes		_				
Have you actually had any thoughts of killing yourself?								
If yes, describe:								
3. Active Suicidal Ideation with Any Methods (Not Plan) without Intent to Act								
Subject endorses thoughts of suicide and has thought of at least one method during the assessment period. This is different than a			No	Yes	No			
specific plan with time, place or method details worked out (e.g. thought of method to kill self but not a specific plan). Includes person who would say, "I thought about taking an overdose but I never made a specific plan as to when, where or how I would actually do								
itand I would never go through with it."	a specific plan as to when, where or now I would actually ao							
Have you been thinking about how you might do this?								
If yes, describe:								
4. Active Suicidal Ideation with Some Intent to Act, with	out Specific Plan							
Active suicidal thoughts of killing oneself and subject reports having so		Yes	No	Yes	No			
thoughts but I definitely will not do anything about them."								
Have you had these thoughts and had some intention of acting on the	m?							
If yes, describe:								
5. Active Suicidal Ideation with Specific Plan and Intent								
Thoughts of killing oneself with details of plan fully or partially worked		Yes	No	Yes	No			
Have you started to work out or worked out the details of how to kill y	ourseig? Do you intena to carry out this plan?							
If yes, describe:								
INTENSITY OF IDEATION								
The following features should be rated with respect to the most								
the least severe and 5 being the most severe). Ask about time h	e/she was feeling the most suicidal.							
<u>Lifetime</u> - Most Severe Ideation:			Most		Most			
Type # (1-5) Description of Ideation			ere	Severe				
Past X Months - Most Severe Ideation:								
Type # (1-5)	Description of Ideation							
Frequency How many times have you had these thoughts?								
How many times have you had these thoughts? (1) Less than once a week (2) Once a week (3) 2-5 times in week (4) Daily or almost daily (5) Many times each day								
Duration								
When you have the thoughts how long do they last?								
(1) Fleeting - few seconds or minutes (2) Less than 1 hour/some of the time	(4) 4-8 hours/most of day (5) More than 8 hours/persistent or continuous							
(3) 1-4 hours/a lot of time	(5) More than 6 nours, persistent of continuous							
Controllability								
Could/can you stop thinking about killing yourself or want (1) Easily able to control thoughts	ting to die if you want to? (4) Can control thoughts with a lot of difficulty							
(2) Can control thoughts with little difficulty	(5) Unable to control thoughts							
(3) Can control thoughts with some difficulty	(0) Does not attempt to control thoughts							
Deterrents Are there things - anyone or anything (e.g., family, religion	n nain of death) - that stopped you from wanting to							
die or acting on thoughts of committing suicide?	i, pain of acain, that stopped you from waiting to							
(1) Deterrents definitely stopped you from attempting suicide	(4) Deterrents most likely did not stop you							
(2) Deterrents probably stopped you(3) Uncertain that deterrents stopped you	(5) Deterrents definitely did not stop you(0) Does not apply							
Reasons for Ideation	(o) Book not apply							
What sort of reasons did you have for thinking about wanting to die or killing yourself? Was it to end the pain								
or stop the way you were feeling (in other words you couldn't go on living with this pain or how you were								
feeling) or was it to get attention, revenge or a reaction from others? Or both? (1) Completely to get attention, revenge or a reaction from others (4) Mostly to end or stop the pain (you couldn't go on					_			
(2) Mostly to get attention, revenge or a reaction from others living with the pain or how you were feeling)								
(3) Equally to get attention, revenge or a reaction from others and to end/stop the pain	living with the pain or how you were feeling) (5) Completely to end or stop the pain (you couldn't go on living with the pain or how you were feeling)							

CIDAL BEHAVIOR cck all that apply, so long as these are separate events; must ask about all types)			Lifetime		Past Years	
Actual Attempt:		Yes	No	Yes	No	
A potentially self-injurious act committed with at least some wish to die, as a result of act. Behavior was in part thought of as method to kill oneself. Intent does not have to be 100%. If there is any intent/desire to die associated with the act, then it can be considered an actual suicide attempt. There does not have to be any injury or harm , just the potential for injury or harm. If person pulls trigger while gun is in mouth but gun is broken so no injury results, this is considered an attempt. Inferring Intent: Even if an individual denies intent/wish to die, it may be inferred clinically from the behavior or circumstances. For example, a highly lethal act that is clearly not an accident so no other intent but suicide can be inferred (e.g., gunshot to head, jumping from window of a high floor/story). Also, if someone denies intent to die, but they thought that what they did could be lethal, intent may be inferred.						
Have you made a suicide attempt?		T. 4	1 // C	Tota	1 # of	
Have you done anything to harm yourself?			l # of mpts	Total # of Attempts		
Have you done anything dangerous where you could have died? What did you do? Did you as a way to end your life?						
Did you want to die (even a little) when you? Were you trying to end your life when you?						
Or Did you think it was possible you could have died from?						
Or did you do it purely for other reasons / without ANY intention of killing yourself (like to relieve stress,	feel better,					
get sympathy, or get something else to happen)? (Self-Injurious Behavior without suicidal intent)	,					
If yes, describe:		Yes	No	Yes	No	
Has subject anguard in New Cuisidal Calf Injunious Debaution?						
Has subject engaged in Non-Suicidal Self-Injurious Behavior? Interrupted Attempt:		Yes	No	Yes	No	
When the person is interrupted (by an outside circumstance) from starting the potentially self-injurious act (if not for that, actual attempt would have occurred).						
Overdose: Person has pills in hand but is stopped from ingesting. Once they ingest any pills, this becomes an attempt rather than an interrupted attempt. Shooting: Person has gun pointed toward self, gun is taken away by someone else, or is somehow prevented from pulling trigger. Once they pull the trigger, even if the gun fails to fire, it is an attempt. Jumping: Person is poised to jump, is grabbed and taken down from ledge. Hanging: Person has noose around neck but has not yet started to hang - is stopped from doing so.				T	1 11 C	
Has there been a time when you started to do something to end your life but someone or something stopped you before					ıl # of rupted	
you actually did anything? If yes, describe:				_		
Aborted Attempt:		Yes	No	Yes	No	
When person begins to take steps toward making a suicide attempt, but stops themselves before they actually have engaged in any self-destructive behavior. Examples are similar to interrupted attempts, except that the individual stops him/herself, instead of being stopped by something else.						
Has there been a time when you started to do something to try to end your life but you stopped yourself before you actually did anything? If yes, describe:					Total # of aborted	
Preparatory Acts or Behavior: Acts or preparation towards imminently making a suicide attempt. This can include anything beyond a verbalization or thought, such as assembling a specific method (e.g., buying pills, purchasing a gun) or preparing for one's death by suicide (e.g., giving things away, writing a suicide note). Have you taken any steps towards making a suicide attempt or preparing to kill yourself (such as collecting pills, getting a gun, giving valuables away or writing a suicide note)? If yes, describe:			No	Yes	No	
Suicidal Behavior:			No	Yes	No	
Suicidal behavior was present during the assessment period?						
		Most Leth	nal	Initial/F		
	•	Attempt Date:		Attempt Date:		
Actual Lethality/Medical Damage:	Enter Code		ode		Code	
 No physical damage or very minor physical damage (e.g., surface scratches). Minor physical damage (e.g., lethargic speech; first-degree burns; mild bleeding; sprains). Moderate physical damage; medical attention needed (e.g., conscious but sleepy, somewhat responsive; second-degree burns; bleeding of major vessel). Moderately severe physical damage; medical hospitalization and likely intensive care required (e.g., comatose with reflexes intact; third-degree burns less than 20% of body; extensive blood loss but can recover; major fractures). Severe physical damage; medical hospitalization with intensive care required (e.g., comatose without reflexes; third-degree burns over 20% of body; extensive blood loss with unstable vital signs; major damage to a vital area). Death 		Enter C	Enter Code Enter Cod			
al Lethality: Only Answer if Actual Lethality=0 lethality of actual attempt if no medical damage (the following examples, while having no actual medical damage, had all for very serious lethality: put gun in mouth and pulled the trigger but gun fails to fire so no medical damage; laying tracks with oncoming train but pulled away before run over).		Enter C	Code	Enter Code		
0 = Behavior not likely to result in injury 1 = Behavior likely to result in injury but not likely to cause death 2 = Behavior likely to result in death despite available medical care						